



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

207. Proposed by F. P. MATZ, Sc. D., Ph. D.

If K represents the complete elliptic integral of the first kind, prove that

$$\int_0^1 \frac{Kd\kappa}{1+\kappa} = \frac{1}{4}\pi^2.$$

208. Proposed by F. P. MATZ, Sc. D., Ph. D.

Solve the differential equation

$$(a^2 + x^2) \frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} = 0.$$

MECHANICS.

185. Proposed by J. EDWARD SANDERS.

A perfectly flexible rope whose weight is w per linear unit, and length $2l$, rests in equilibrium on a smooth peg. If now one end be raised a distance a and then released, find the time in which this end will rise to the height x above its original position, and the tension at that instant of the rope at the point where it passes over the peg.

MISCELLANEOUS.

152. Proposed by J. EDWARD SANDERS.

A conductor, the equation of the surface of which is

$$\frac{x^2}{25} + \frac{y^2}{16} + \frac{z^2}{9} = 1,$$

is charged with 80 units of electricity, what is the density at a point for which $x=3$, $y=3$? If the density of this point be a , what is the whole charge on the ellipsoid? [From Peirce's *Potential Functions*, example 165, p. 388.]

153. Proposed by CHRISTIAN HORNING, A. M., Heidelberg University, Tiffin, Ohio.

Two men start from Columbus, Ohio, at the same time; one travels east and the other west. They travel at the rate of 4 miles an hour from sunrise to sunset each day until they meet. Where will they meet and what distance will each have traveled?

154. Proposed by D. BIDDLE (Unsolved problem in the Educational Times, London).

Prove that the proper angle at which to cross a street when a person wishes to continue his course on the other side, and the roadway is n times as muddy as the pavement, is that of which the sine is $(n^2 - 1)/(n^2 + 1)$.